

DFPD 97/TH/22
May 1997

A FERMIPHOBIC $SU(2) \times SU(2) \times U(1)$ EXTENSION OF THE SM

J. Matias

Dipartimento di Fisica.

Università di Padova.

Via F. Marzolo 8, I-35131 Padova, Italy

Abstract

A new type of gauge extension of the SM is proposed. It is based on a $SU(2) \times SU(2) \times U(1)$ group with the peculiarity that the gauge bosons of the extra $SU(2)$ do not couple to fermions. This feature relaxes some of the constraints on the masses of the new gauge bosons, leaving the possibility of having lighter masses than in traditional extensions. The model exhibits several interesting properties, it is anomaly free and at tree level it does not have FCNC while loop induced effects are strongly suppressed. Also, from the analysis of $\Delta\rho$ at one loop, two configurations for the vevs giving rise to a screening phenomenon are identified. One of these configurations can also be related with the Bess model. A tree level fit to the most recent electroweak data is performed confirming the possibility of having new light gauge boson masses. The constraints coming from different FCNC processes like $b \rightarrow s\gamma$, $B_0 - \bar{B}_0$ and $K_0 - \bar{K}_0$ mixing are also taken into account. Finally, a generalization of this model for the case of having several extra copies of $SU(2)$ groups is commented, focusing on the presence of screening configurations and the corresponding mass spectrum.

Contribution to the XXXIIInd Rencontres de Moriond:
"Electroweak Interactions and Unified Theories", Les Arcs, France, March 1997

In the post-LEPI era the extraordinary success of the Standard Model (SM) in explaining data has been confirmed. As a consequence any acceptable alternative should be able to repeat its success while allowing for new physics at a not very high scale to be interesting for present and future colliders. Extensions of the SM based on enlarging the gauge group with an extra $SU(2)$ could fulfill this requirement. However, in most of these models (Standard Left-Right (LR)¹⁾ or ununified models²⁾) the lower bounds on the new gauge boson masses are still quite high and new problems like anomalies or Flavour Changing Neutral Currents (FCNC) appear.

Instead, one can think of a particular type of $SU(2)$ extension of the SM that we have called ‘fermiophobic’³⁾, which seems to naturally satisfy all existing constraints and, at the same time, avoids most of the problems of the other models while allowing for light new gauge bosons.

This model, defined by the transformation properties of fermions under the gauge group $SU(2)_L \times SU(2)_R \times U(1)_{\tilde{Y}}$,

$$\begin{aligned} q_L &= \begin{pmatrix} u_L \\ d_L \end{pmatrix} \sim (2, 1, 1/6) & l_L &= \begin{pmatrix} \nu_L \\ e_L \end{pmatrix} \sim (2, 1, -1/2) \\ u_R &\sim (1, 1, 2/3) & d_R &\sim (1, 1, -1/3) & e_R &\sim (1, 1, -1) \end{aligned}$$

is automatically free of gauge anomalies. Contrary to standard LR models, it does not require the existence of right-handed neutrinos.

The field content of the scalar sector consists of two doublets and one bidoublet

$$\phi_L = \begin{pmatrix} \phi_L^0 \\ \phi_L^- \end{pmatrix} \sim (2, 1, -1/2) \quad \phi_R = \begin{pmatrix} \phi_R^0 \\ \phi_R^- \end{pmatrix} \sim (1, 2, -1/2) \quad \phi_{LR} = \begin{pmatrix} \phi_1^0 & \phi_2^+ \\ \phi_1^- & \phi_2^0 \end{pmatrix} \sim (2, 2, 0)$$

The absence of right-handed neutrinos makes unnecessary the use of scalar triplets. If we assume that there are no new sources of CP-violation, the previous sixteen scalar degrees of freedom organize in a set of eight charged scalars: four Goldstone bosons ($G^\pm, G'^\pm$) and four charged physical scalars $H_1^\pm, H_2^\pm$ with mixing angle $\beta_\pm$. And eight neutrals: four scalars (H_3^0, H_4^0, H_5^0 and H_6^0), one of them can be identified with the SM-Higgs like, and four pseudoscalars, two Goldstone bosons G^0, G'^0 and two physical pseudoscalars H_1^0, H_2^0 with mixing angle β_0 .

Finally, concerning the gauge sector we have four charged gauge bosons ($W^\pm, W'^\pm$) with mixing angle $\alpha_\pm$, the photon γ and two neutrals (Z, Z') with mixing angle α_0 . Moreover, the gauge coupling constants g_L and g_R can be different and its ratio g_R/g_L is called x .

Given these quantum number assignments and assuming a pattern of real vevs for the scalar fields

$$\langle \phi_L \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} v_L \\ 0 \end{pmatrix} \quad \langle \phi_R \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} v_R \\ 0 \end{pmatrix} \quad \langle \phi_{LR} \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} v_1 & 0 \\ 0 & v_2 \end{pmatrix},$$

one can automatically write down the lagrangian and derive the gauge boson masses (both doublets and the bidoublet contribute) and the relevant vertices³⁾. Notice that the Yukawa part of the lagrangian turns out to be as in the SM, only the ϕ_L doublet can couple to fermions, avoiding automatically the tree level FCNC problems of standard LR

models. Moreover, at the level of gauge interactions of fermions an important remark is the strong suppression of the $W'^{\pm}$ (mass eigenstate) coupling to fermions, while the corresponding Z' coupling is suppressed only in the large g_R limit.

We have imposed a theoretical constraint to the model at the level of radiative corrections. We require that all contributions quadratic in the masses of the scalars cancel in $\Delta\rho$ at one-loop. The implications of such *screening* requirement however, go well beyond to the mere cancellation of quadratic contributions. From the explicit expression of $\Delta\rho$ ⁴⁾ one sees that the fulfilment of that condition can be reached in different ways. Two of them translate into constraints on the vevs, β angles and scalar masses:

i) $v_R \rightarrow \infty$ and $\beta_{\pm} = \beta_0$. ρ at tree level is exactly one and the one-loop correction corresponds to the one of a two doublet model. However, the new gauge boson masses become infinite.

ii) $v_R = v_L/x$, $v_1 = v_2 = v$, $\beta_{\pm} = \beta_0$ and $\tan\beta = -M_{W'}/M_W$. When the mass of the H_1^+ is degenerate with H_5^0 and H_2^+ with H_2^0 the one-loop quadratic contributions to $\Delta\rho$ cancel. Masses and mixing angles become extremely simple in that case,

$$M_W^2 = \frac{g_L^2 v_L^2}{4} \quad M_Z^2 = \frac{g_L^2 v_L^2}{4} \frac{x^2}{x^2 - s_W^2 x^2 - s_W^2} \quad M_{W'}^2 = M_{Z'}^2 = \frac{g_L^2}{4} (v_L^2 + 2v^2(1 + x^2))$$

and the mixing angles are $\tan\alpha_{\pm} = 1/x$ and $\tan\alpha_0 = 1/\sqrt{x^2 - s_W^2 x^2 - s_W^2}$ with s_W the sinus of the Weinberg angle. Notice from the previous expressions that the new gauge bosons should be degenerate with a mass always heavier than M_W . One observes, also, that the theory requires x to be large in order to have a small tree level ρ parameter. Moreover, an interesting property of that configuration is that when one imposes a further restriction on the vevs ($v_R = \sqrt{2}v$) the mass predictions of our model coincide with those of the Bess model⁵⁾ in a restricted 4-parameter space.

One of the most important experimental constraints on the model comes from a tree level fit to the electroweak data at the Z peak, the low energy data from neutrino-hadron scattering and atomic parity violation experiments (fourteen observables in total)⁶⁾. The model has, concerning only bosonic interactions, seven parameters (g_L , g_R , $\tilde{g}$, v_R , v_L , v_1 , v_2) that we have translated in terms of the three input quantities of the SM precision tests (G_F , α , M_Z) and four extra input parameters: $\alpha_{\pm}$, α_0 , $M_{W'}$ and x . The fit was done by adding to the SM predictions the deviations due to the model. These deviations depend on the four extra input parameters. A χ^2 minimization procedure is used to determine their best values.

The results of the fit are:

a) in the general case the fit turns out to be quite insensitive to $M_{W'}$ for fixed values of x . So we kept x and $M_{W'}$ fixed in a range $1 \leq x \leq 15$ and $100 \leq M_{W'} \leq 1000$ leaving $\alpha_{\pm}$ and α_0 free. In that scenario the experimental data allows a mixing angle $\alpha_{\pm}, \alpha_0$ in the range $10^{-2} - 10^{-3}$ and larger mixing angles are allowed for larger x values. The masses of the new gauge bosons W' , Z' can be in a broad range allowing also for light values, even of the order of 150 GeV if x is sufficiently large ($x > 5$).

b) in the particular screening configuration ii) mentioned above, we no more have four but just two free parameters (x and $M_{W'}$). The fit prefers large values of x approaching in that limit the SM and becoming then insensitive to the value of $M_{W'}$.

Other bounds on the masses of the new gauge bosons and the charged Higgs bosons could come from one-loop FCNC processes. While they are quite severe in standard LR models⁷⁾, in our model they become naturally relaxed due to the absence of coupling between the ϕ_R , the ϕ_{LR} and the $W_R^\pm$ field (in the interacting basis) with ordinary fermions. The $b \rightarrow s\gamma$, $B_0 - \bar{B}_0$ and $K_0 - \bar{K}_0$ mixing have been examined finding an exclusion region for light values of the mass of the charged Higgs bosons (we take them to be degenerate) when small values of x are taken. However, no sensitivity to $M_{W'}$ was found. In the $b \rightarrow s\gamma$ process, for instance, this is due to the strong suppression of the first correction to the $W^\pm$ boson exchange amplitude that, in our model, is of order $\alpha_\pm^2$ while in standard LR models is of order $\alpha_\pm$.

Bounds like those coming from neutrinoless double beta decay are, obviously, innocuous to our model.

The study of the limits from the Tevatron, and, of the possible signals of the new gauge boson particles in present and future colliders is presently being completed³⁾.

Finally the existence of such screening configurations has been studied also in a more general case with additional $SU(2)$ groups⁴⁾. The result, again positive, is particularly smart since it allows to obtain the exact gauge mass spectrum of an $SU(2) \times SU(2)^n \times U(1)$ theory starting just from the information of the screening of quadratic contributions to $\Delta\rho$. The pattern of new gauge boson masses is similar to that of the case ii). Each pair of new gauge bosons masses is degenerate and differs from the other pairs. M_W is exactly the same as in the case $n = 1$ and the M_Z mass is a straightforward generalization of the $n = 1$ case.

The work reported here was done in collaboration with A. Donini, F. Feruglio and F. Zwirner.

References

- [1] J.C. PATI AND A. SALAM, *Phys. Rev.* **D10** (1974) 275; R.N. MOHAPATRA AND J.C. PATI, *Phys. Rev.* **D11** (1975) 566 and 2558; G. SENJANOVIC AND R.N. MOHAPATRA, *Phys. Rev.* **D12** (1975) 1502.
- [2] H. GEORGI, E.E. JENKINS AND E.H. SIMMONS, *Nucl. Phys.* **B331** (1990) 541.
- [3] A. DONINI, F. FERUGLIO, J. MATIAS AND F. ZWIRNER, "Phenomenological aspects of a fermiophobic $SU(2) \times SU(2) \times U(1)$ extension of the SM", Padova/Cern Prep. DFPD 97/TH/23 1997.
- [4] J. MATIAS AND A. VICINI, "The ρ parameter and the screening phenomena for sequential W and Z gauge bosons", Padova/Desy Prep. 1997.
- [5] R. CASALBUONI, S. DE CURTIS, D. DOMINICI AND R. GATTO, *Nucl. Phys.* **B282** (1987) 235.
- [6] PARTICLE DATA GROUP, *Phys. Rev.* **D54** (1996) 1; The LEP Electroweak Working Group; prep. CERN-PPE/96-183 and Internal Note LEPEWWG 97-01; C.S. WOOD ET AL. *Science* **275** (1997) 1759.
- [7] P. LANGACKER AND S. UMA SANKAR, *Phys. Rev.* **D40** (1989) 1569.